

## Two-Colorings of Positive Integers

STEVEN FINCH

May 27, 2008

Let  $f : \{1, 2, 3, \dots\} \rightarrow \{-1, 1\}$  be an arbitrary function. Given a threshold  $M > 0$ , we ask two questions:

- Do there exist integers  $a > 0$ ,  $b \geq 0$ ,  $\ell > 0$  such that

$$|f(a + b) + f(2a + b) + f(3a + b) + \dots + f(\ell a + b)| > M?$$

- Do there exist integers  $a > 0$ ,  $\ell > 0$  such that

$$|f(a) + f(2a) + f(3a) + \dots + f(\ell a)| > M?$$

The answer to the first question is yes. In words, every two-coloring of the positive integers has unbounded discrepancy, taken over the family of arithmetic progressions. Restricting attention to the subset  $\{1, 2, 3, \dots, n\}$ , we have [1, 2, 3, 4]

$$c n^{1/4} \leq P(n) = \min_f \max_{\substack{a, b, \ell \\ \ell a + b \leq n}} \left| \sum_{k=1}^{\ell} f(k a + b) \right| \leq C n^{1/4}$$

for all  $n$ , with constants  $c \geq 1/20$  and  $C < \infty$ . The lower bound on  $c$  is improved to  $1/7$  in [5]; no finite upper bound on  $C$  is apparently known. It is natural to wonder about the numerical values of

$$\liminf_{n \rightarrow \infty} n^{-1/4} P(n), \quad \limsup_{n \rightarrow \infty} n^{-1/4} P(n).$$

The second question, due to Erdős [6, 7, 8] and Chudakov [9, 10], remains open. It is remarkable that, upon mere constraint to homogeneity ( $b = 0$ ), the problem becomes unsolved! If we expand the family under consideration, more can be said. For almost all real numbers  $\alpha \geq 1$ , there exists  $\ell > 0$  such that [11, 12, 13]

$$|f(\lfloor \alpha \rfloor) + f(\lfloor 2\alpha \rfloor) + f(\lfloor 3\alpha \rfloor) + \dots + f(\lfloor \ell \alpha \rfloor)| > M.$$

---

<sup>0</sup>Copyright © 2008 by Steven R. Finch. All rights reserved.

Such **quasi-arithmetic progressions** collapse to homogeneous arithmetic progressions when  $\alpha$  is an integer. Even though the set  $S$  of counterexamples  $\alpha$  has measure zero, it is not known whether  $S$  avoids all integers.

We also examine the expression [14, 15]

$$Q(n) = \min_f \max_{\substack{a,b,\ell \\ a < b \\ \ell + b \leq n}} \left| \sum_{k=1}^{\ell} f(k+a)f(k+b) \right|$$

and wonder about the numerical values of

$$\liminf_{n \rightarrow \infty} n^{-1/2} Q(n), \quad \limsup_{n \rightarrow \infty} n^{-1/2} Q(n).$$

**0.1. Addendum.** If  $f$  is random (independently taking the values  $\pm 1$  with probability  $1/2$  at each integer  $1 \leq k \leq n$ ), then an asymptotic statement can be made about the mean: [16]

$$\mathbb{E}(|f(1) + f(2) + f(3) + \cdots + f(n)|) \sim \sqrt{\frac{2n}{\pi}}$$

as  $n \rightarrow \infty$ , and likewise

$$\mathbb{E}(|f(1)f(1+b) + f(2)f(2+b) + f(3)f(3+b) + \cdots + f(n-b)f(n)|) \sim \sqrt{\frac{2n}{\pi}}$$

for fixed  $b \geq 1$ . A proof of the latter can be based on [17]; the order of  $Q(n)$  is  $n^{1/2}$  (in agreement) whereas the order of  $P(n)$  is only  $n^{1/4}$  (in disagreement).

Via [18, 19], we learned of the Polymath wiki – which documents massively collaborative online mathematical projects – and which includes work on problems given here [20].

#### REFERENCES

- [1] K. F. Roth, Remark concerning integer sequences, *Acta Arith.* 9 (1964) 257–260; MR0168545 (29 #5806).
- [2] J. Beck, Roth’s estimate of the discrepancy of integer sequences is nearly sharp, *Combinatorica* 1 (1981) 319–325; MR0647981 (83i:05040).
- [3] J. Beck and V. T. Sós, Discrepancy theory, *Handbook of Combinatorics*, v. 2, ed. R. L. Graham, M. Grötschel and L. Lovász, MIT Press, 1995, pp. 1405–1446; MR1373682 (96m:11060).

- [4] J. Matoušek and J. Spencer, Discrepancy in arithmetic progressions, *J. Amer. Math. Soc.* 9 (1996) 195–204; MR1311824 (96c:11089).
- [5] L. Lovász, Integer sequences and semidefinite programming, *Publ. Math. Debrecen* 56 (2000) 475–479; MR1765994 (2001k:11020).
- [6] P. Erdős, Quelques problèmes de théorie des nombres, *Monographies de L'Enseignement Mathématique*, v. 6, L'Enseignement Mathématique, Université, Geneva, 1963, pp. 81–135; MR0158847 (28 #2070).
- [7] P. Erdős, Some unsolved problems, *Michigan Math. J.* 4 (1957) 291–300; MR0098702 (20 #5157).
- [8] P. Erdős, On the combinatorial problems which I would most like to see solved, *Combinatorica* 1 (1981) 25–42; MR0602413 (82k:05001).
- [9] N. G. Tchudakoff, Theory of the characters of number semigroups, *J. Indian Math. Soc.* 20 (1956) 11–15; MR0083515 (18,719e).
- [10] A. R. D. Mathias, On a conjecture of Erdős and Čudakov, *Combinatorics, Geometry and Probability*, Proc. 1993 Cambridge conf., ed. B. Bollobás and A. Thomason, Cambridge Univ. Press, 1997, pp. 487–488; MR1476466 (98i:11058).
- [11] J. Beck, On irregularities of  $\pm 1$ -sequences, *Österreichische Akademie der Wissenschaften Mathematisch-Naturwissenschaftliche Klasse. Sitzungsberichte. Abteilung II* 195 (1986) 13–23; MR0881327 (88e:11065).
- [12] R. Hochberg, Large discrepancy in homogeneous quasi-arithmetic progressions, *Combinatorica* 26 (2006) 47–64; MR2201283 (2006i:11012).
- [13] S. Vijay, On the discrepancy of quasi-progressions, math.CO/0604511.
- [14] J. Cassaigne, C. Mauduit and A. Sárközy, On finite pseudorandom binary sequences. VII, The measures of pseudorandomness, *Acta Arith.* 103 (2002) 97–118; MR1904866 (2004c:11139).
- [15] N. Alon, Y. Kohayakawa, C. Mauduit, C. G. Moreira and V. Rödl, Measures of pseudorandomness for finite sequences: minimal values, *Combin. Probab. Comput.* 15 (2006) 1–29; MR2195573 (2006j:60007).
- [16] S. R. Finch, Moments of sums, unpublished note (2004).
- [17] H. C. Ho and T. C. Sun, A central limit theorem for noninstantaneous filters of a stationary Gaussian process, *J. Multivariate Anal.* 22 (1987) 144–155; MR0890889 (88m:60059).

- [18] M. Coons, On the multiplicative Erdős discrepancy problem, arXiv:1003.5388.
- [19] J. Vandehey, On multiplicative functions with bounded partial sums, arXiv:1108.1487.
- [20] The Erdős discrepancy problem, <http://michaelnielsen.org/polymath1>.