

Lane-Ritter-Emden Constants

STEVEN FINCH

November 30, 2008

The Lane-Emden equation of index p :

$$y''(x) + \frac{2}{x}y'(x) + y(x)^p = 0, \quad y(0) = 1, \quad y'(0) = 0$$

is useful in astrophysics for computing the structure of interiors of polytropic stars [1, 2, 3]. A well-known series for $y(x)$ is [4]

$$y(x) = \sum_{k=0}^{\infty} a_k x^{2k}, \quad x \approx 0$$

where $a_0 = 1$, $a_1 = -1/6$ and

$$a_k = \frac{1}{(k-1)k(2k+1)} \sum_{j=1}^{k-1} (jp + j - k + 1)(k-j)(2k-2j+1)a_j a_{k-j}, \quad k \geq 2.$$

This series has radius of convergence [5, 6, 7, 8, 9, 10]

$$\gamma = \lim_{k \rightarrow \infty} \left| \frac{a_k}{a_{k+1}} \right|^{1/2} = \begin{cases} \infty & \text{if } p = 0 \text{ or } p = 1, \\ 3.6537537362\dots = \sqrt{13.3499163649\dots} & \text{if } p = 3/2, \\ 3.9645856345\dots = \sqrt{15.7179392534\dots} & \text{if } p = 2, \\ 2.5748367419\dots = \sqrt{6.6297842476\dots} & \text{if } p = 3, \\ 2.0348941557\dots = \sqrt{4.1407942251\dots} & \text{if } p = 4, \\ 1.7320508075\dots = \sqrt{3} & \text{if } p = 5 \end{cases}$$

and details on relevant calculations will appear momentarily. The *dimensionless radius* of a polytropic star is the smallest positive x_0 for which $y(x_0) = 0$: [1, 11, 12, 13, 14, 15]

$$x_0 = \begin{cases} 2.4494897427\dots = \sqrt{6} & \text{if } p = 0, \\ 3.1415926535\dots = \pi & \text{if } p = 1, \\ 3.6537537362\dots = \sqrt{13.3499163649\dots} & \text{if } p = 3/2, \\ 4.3528745959\dots = \sqrt{18.9475172480\dots} & \text{if } p = 2, \\ 6.8968486193\dots = \sqrt{47.5665208786\dots} & \text{if } p = 3, \\ 14.9715463488\dots = \sqrt{224.1472000754\dots} & \text{if } p = 4, \\ \infty & \text{if } p = 5 \end{cases}$$

⁰Copyright © 2008 by Steven R. Finch. All rights reserved.

and the *dimensionless mass* of a polytropic star is x_0^2 multiplied by $-y'(x_0)$:

$$\mu = -x_0^2 y'(x_0) = \begin{cases} 4.8989794855\dots = 2\sqrt{6} & \text{if } p = 0, \\ 3.1415926535\dots = \pi & \text{if } p = 1, \\ 2.7140551201\dots & \text{if } p = 3/2, \\ 2.4110460120\dots & \text{if } p = 2, \\ 2.0182359509\dots & \text{if } p = 3, \\ 1.7972299144\dots & \text{if } p = 4, \\ 1.7320508075\dots = \sqrt{3} & \text{if } p = 5. \end{cases}$$

No closed-form expressions for constants associated with the range $1 < p < 5$ are known. The functions $\gamma(p)$ and $x_0(p)$ are initially equal for $p > 1$, but they separate at $p \approx 1.9121$ [7]. The function $\mu(p)$ initially decreases, but encounters a minimum at $p \approx 4.823$ and increases henceforth [16, 17].

A simpler formula for the coefficients $\{a_k\}$ is valid for $p = 2$: [18]

$$a_k = \frac{-1}{(2k)(2k+1)} \sum_{j=0}^{k-1} a_j a_{k-j}, \quad k \geq 1$$

which makes the alternating character of the series obvious. Is there an analogous formula for $p = 3$ or $p = 4$?

Let us explain how $\gamma(p)$ is computed for $2 \leq p \leq 5$. Write $t = -x^2$ and $u = y^{-1/p}$, then

$$\begin{aligned} -6p u \frac{du}{dt} + 4p(p+1)t \left(\frac{du}{dt} \right)^2 - 4p t u \frac{d^2 u}{dt^2} &= u^{-p^2+p+2} \\ u(0) = 1, \quad \frac{du}{dt}(0) &= -\frac{1}{6p}. \end{aligned}$$

For example, supposing $p = 2$, we find $u(10) = 0.312\dots$ and $\frac{du}{dt}(10) = -0.058\dots$. By the Inverse Function Theorem,

$$-6p u \left(\frac{dt}{du} \right)^2 + 4p(p+1)t \frac{dt}{du} + 4p t u \frac{d^2 t}{du^2} = u^{-p^2+p+2} \left(\frac{dt}{du} \right)^3.$$

In the case $p = 2$, initial conditions $t(0.312\dots) = 10$ and $\frac{dt}{du}(0.312\dots) = \frac{1}{-0.058\dots}$ clearly hold. We find $t(0) = 15.717\dots$, thus $x = (\pm 3.964\dots)i$ correspond to where $y = u^{-p}$ explodes [19]. This technique works because 10 is large enough that $u(10)$ is small, making the computation of $t(0)$ feasible.

See also [26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62]; we hope to report on [63, 64, 65, 66, 67, 68] later.

0.1. Polytropic and Isothermal Spheres. A generalization of the Lane-Emden equation is [17, 20, 21, 22, 23, 24]

$$y''(x) + \frac{N}{x}y'(x) + y(x)^p = 0, \quad y(0) = 1, \quad y'(0) = 0$$

corresponding to N -dimensional polytropic spheres in \mathbb{R}^{N+1} . The case $N = 2$ was discussed earlier. For $N = 1$ (polytropic cylinders), we have

$$x_0 = \begin{cases} 2 & \text{if } p = 0, \\ 2.4048255576\dots = z & \text{if } p = 1, \\ 2.6477767662\dots & \text{if } p = 3/2, \\ 2.9213207237\dots & \text{if } p = 2, \\ 3.5739009819\dots & \text{if } p = 3 \end{cases}$$

where z is the smallest positive zero of the Bessel function J_0 and [25]

$$\mu = -x_0 y'(x_0) = \begin{cases} 2 & \text{if } p = 0, \\ 1.2484591696\dots = z J_1(z) & \text{if } p = 1, \\ 1.0611147888\dots & \text{if } p = 3/2, \\ 0.9253532703\dots & \text{if } p = 2, \\ 0.7401221205\dots & \text{if } p = 3. \end{cases}$$

No closed-form expressions for constants associated with $p > 1$ are known. In contrast, for $N = 0$ (polytropic slabs),

$$x_0 = \left(\frac{\pi}{2(p+1)} \right)^{1/2} \frac{\Gamma\left(\frac{1}{p+1}\right)}{\Gamma\left(\frac{p+3}{2(p+1)}\right)} = \begin{cases} 1.4142135623\dots = \sqrt{2} & \text{if } p = 0, \\ 1.5707963267\dots = \pi/2 & \text{if } p = 1, \\ 1.6453408471\dots & \text{if } p = 3/2, \\ 1.7173153422\dots & \text{if } p = 2, \\ 1.8540746773\dots & \text{if } p = 3 \end{cases}$$

and

$$\mu = -y'(x_0) = \left(\frac{2}{p+1} \right)^{1/2} = \begin{cases} 1.4142135623\dots = \sqrt{2} & \text{if } p = 0, \\ 1 & \text{if } p = 1, \\ 0.8944271909\dots & \text{if } p = 3/2, \\ 0.8164965809\dots & \text{if } p = 2, \\ 0.7071067811\dots = 1/\sqrt{2} & \text{if } p = 3. \end{cases}$$

A different generalization involves the limit as $p \rightarrow \infty$:

$$y''(x) + \frac{2}{x}y'(x) = e^{-y(x)}, \quad y(0) = y'(0) = 0.$$

This corresponds to 2-dimensional isothermal spheres in \mathbb{R}^3 and has the following series expansion:

$$y(x) = \sum_{k=1}^{\infty} b_k x^{2k}, \quad x \approx 0$$

where $b_1 = 1/6$ and

$$b_k = \frac{-1}{(k-1)k(2k+1)} \sum_{j=1}^{k-1} j(k-j)(2k-2j+1)b_j b_{k-j}, \quad k \geq 2.$$

The radius of convergence, squared, is [7, 8, 69]

$$\lim_{k \rightarrow \infty} \left| \frac{b_k}{b_{k+1}} \right| = 10.7170288238\dots = 2(5.3585144119\dots).$$

This is computed by writing $t = -x^2$ and $u = e^{y/2}$, then applying the Inverse Function Theorem to

$$\begin{aligned} -12u \frac{du}{dt} + 8t \left(\frac{du}{dt} \right)^2 - 8t u \frac{d^2u}{dt^2} &= 1, \\ u(0) = 1, \quad \frac{du}{dt}(0) &= -\frac{1}{12}. \end{aligned}$$

It is also known that

$$-y(x) \sim \ln \left(\frac{2}{x^2} \right) + \frac{C}{\sqrt{x}} \cos \left(\frac{\sqrt{7}}{2} \ln(x) - c \right)$$

as $x \rightarrow \infty$ for certain unspecified constants C and c . A more precise statement of this asymptotic formula, with expressions for C and c , would be good to see. Related materials include [70, 71, 72, 73, 74, 75].

REFERENCES

- [1] S. Chandrasekhar, *An Introduction to the Study of Stellar Structure*, Dover, 1957, pp. 84–182; MR0092663 (19,1142b).
- [2] M. H. Wrubel, Stellar interiors, *Encyclopedia of Physics*, v. 51, *Astrophysics II: Stellar Structure*, ed. S. Flügge, Springer-Verlag, 1958, pp. 1–74.
- [3] H. T. Davis, *Introduction to Nonlinear Differential and Integral Equations*, Dover, 1962, pp. 371–394, 408; MR0181773 (31 #6000).
- [4] Z. F. Seidov and R. K. Kuzakhmedov, Solution of the Lane-Emden problem in series (in Russian), *Astronomicheskii Zh.* 54 (1977) 707–708; Engl. transl. in *Soviet Astronomy* 21 (1977) 399–400; MR0537843 (80j:85007).

- [5] K. A. Thernöe, Numerische Untersuchungen über die Emden-Funktionen. I, *Z. Astrophys.* 22 (1943) 122–155; MR0009675 (5,182f).
- [6] I. W. Roxburgh and L. M. Stockman, Power series solutions of the polytropic equation, *Monthly Notices Royal Astron. Soc.* 303 (1999) 466–470.
- [7] C. Hunter, Series solutions for polytropes and the isothermal sphere, *Monthly Notices Royal Astron. Soc.* 328 (2001) 839–847.
- [8] A. Aslanov, Determination of convergence intervals of the series solutions of Emden-Fowler equations using polytropes and isothermal spheres, *Phys. Lett. A* 372 (2008) 3555–3561; MR2413857.
- [9] M. I. Nouh, Accelerated power series solution of polytropic and isothermal gas spheres, *New Astronomy* 9 (2004) 467–473.
- [10] A. S. Saad, Symbolic analytical solution of N -dimensional radially symmetric polytropes, *Astron. Nachr.* 325 (2004) 733–739.
- [11] D. H. Sadler, J. C. P. Miller and J. R. Airey, *Mathematical Tables*, v. II, *Emden Functions*, British Assoc. Adv. Sci., 1932.
- [12] C. Mohan and A. R. Al-Bayat, Power series solutions of the Lane-Emden equation, *Astrophys. Space Sci.* 73 (1980) 227–239.
- [13] J. R. Jabbar, The boundary conditions for polytropic gas spheres, *Astrophys. Space Sci.* 100 (1984) 447–449.
- [14] G. P. Horedt, Seven-digit tables of Lane-Emden functions, *Astrophys. Space Sci.* 126 (1986) 357–408.
- [15] Z. F. Seidov, Lane-Emden equation: perturbation method, arXiv:astro-ph/0402130.
- [16] Z. F. Seidov and R. K. Kuzakhmedov, New solutions of the Lane-Emden equation (in Russian), *Astronomicheskii Zh.* 55 (1978) 1250–1255; Engl. transl. in *Soviet Astronomy* 22 (1978) 711–714; MR0537843 (80j:85007).
- [17] G. P. Horedt, Exact solutions of the Lane-Emden equation in N -dimensional space, *Astron. Astrophys.* 160 (1986) 148–156.
- [18] Z. F. Seidov, The power series as solution of the Lane-Emden equation with index two (in Russian), *Dokl. Akad. Nauk AzSSR* 35 (1979) 21–24.

- [19] D. Rusin and R. B. Israel, Convergence radius for a nonlinear ODE solution, unpublished note (2000), http://www.math.niu.edu/~rusin/known-math/00_incoming/emden.
- [20] G. P. Horedt, Approximate analytical solutions of the Lane-Emden equation in N -dimensional space, *Astron. Astrophys.* 172 (1987) 359–367; MR0878810 (88a:85004).
- [21] J. Ostriker, The equilibrium of polytropic and isothermal cylinders, *Astrophys. J.* 140 (1964) 1056–1066; MR0189788 (32 #7208).
- [22] J. Ostriker, Cylindrical Emden and associated functions, *Astrophys. J. Suppl.* 11 (1965) 167–184.
- [23] T. F. Tascione, Plane-symmetric polytropic functions, *Astrophys. J. Suppl.* 24 (1972) 479–492; MR0314224 (47 #2776).
- [24] E. R. Harrison and R. G. Lake, Polytropic and isothermal plane-symmetric configurations, *Astrophys. J.* 171 (1972) 323–330.
- [25] S. R. Finch, Bessel function zeroes, unpublished note (2003).
- [26] R. V. Ramnath, On a class of nonlinear differential equations of astrophysics, *J. Math. Anal. Appl.* 35 (1971) 27–47; MR0277830 (43 #3563).
- [27] Y. Viala and G. P. Horedt, Polytropic sheets, cylinders and spheres with negative index, *Astron. Astrophys.* 33 (1974) 195–202.
- [28] Y. Viala and G. P. Horedt, Emden’s functions for polytropes with negative index, *Astronom. Astrophys. Suppl.* 16 (1974) 173–206; MR0353631 (50 #6114).
- [29] A. P. Linnell, Approximation functions to the Emden and associated Emden functions near the first zero, *Astrophys. Space Sci.* 37 (1975) 57–72.
- [30] A. P. Linnell, Approximation functions to the Emden and associated Emden functions near the first zero. II, *Astrophys. Space Sci.* 37 (1975) 401–426.
- [31] P. Pascual, Lane-Emden equation and Padé approximants, *Astron. Astrophys.* 60 (1977) 161–163.
- [32] A. T. Service, Concise approximation formulae for the Lane-Emden functions, *Astrophys. J.* 211 (1977) 908–909.
- [33] P. G. L. Leach, First integrals for the modified Emden equation $\ddot{q} + \alpha(t)\dot{q} + q^n = 0$, *J. Math. Phys.* 26 (1985) 2510–2514; MR0803793 (86k:34001).

- [34] P. B. Burt, Nonperturbative solutions of the Lane-Emden equation, *Phys. Lett. A* 109 (1985) 133–135; MR0788102 (86h:85001).
- [35] P. B. Burt, Nonperturbative solution of nonlinear field equations, *Nuovo Cimento B* 100 (1987) 43–52; MR0926436 (89h:81021).
- [36] M. Beech, An approximate solution for the polytrope $n = 3$, *Astrophys. Space Sci.* 132 (1987) 393–396.
- [37] C. M. Bender, K. A. Milton, S. S. Pinsky and L. M. Simmons, A new perturbative approach to nonlinear problems, *J. Math. Phys.* 30 (1989) 1447–1455; MR1002247 (90i:34085).
- [38] M. M. Chawla, M. K. Jain and R. Subramanian, The application of explicit Nyström methods to singular second order differential equations, *Comput. Math. Appl.* 19 (1990) 47–51; MR1044440 (91a:65172).
- [39] G. P. Horedt, Analytical and numerical values of Emden-Chandrasekhar associated functions, *Astrophys. J.* 357 (1990) 560–565.
- [40] N. T. Shawagfeh, Nonperturbative approximate solution for Lane-Emden equation, *J. Math. Phys.* 34 (1993) 4364–4369; MR1233277 (94e:85001).
- [41] G. Adomian, R. Rach and N. T. Shawagfeh, On the analytic solution of Lane-Emden equation, *Found. Phys. Lett.* 8 (1995) 161–181.
- [42] B. K. Datta, Analytic solution to the Lane-Emden equation, *Nuovo Cimento B* 111 (1996) 1385–1388; MR1433976 (97j:34019).
- [43] H. Goenner and P. Havas, Exact solutions of the generalized Lane-Emden equation, *J. Math. Phys.* 41 (2000) 7029–7042; MR1781422 (2001d:34004).
- [44] V. B. Mandelzweig and F. Tabakin, Quasilinearization approach to nonlinear problems in physics with application to nonlinear ODEs, *Comput. Phys. Comm.* 141 (2001) 268–281; arXiv:physics/0102041; MR1872036 (2002j:34016).
- [45] H. Goenner, Symmetry transformations for the generalized Lane-Emden equation, *General Relativity and Gravitation* 33 (2001) 833–841.
- [46] Z. F. Seidov, Lane-Emden equation: Picard vs Pade, arXiv:astro-ph/0107395.
- [47] D. C. Biles, M. P. Robinson and J. S. Spraker, A generalization of the Lane-Emden equation, *J. Math. Anal. Appl.* 273 (2002) 654–666; MR1932513 (2003g:34007).

- [48] A.-M. Wazwaz, A new method for solving singular initial value problems in the second-order ordinary differential equations, *Appl. Math. Comput.* 128 (2002) 45–57; MR1891179 (2002m:34011).
- [49] S. Liao, A new analytic algorithm of Lane-Emden type equations, *Appl. Math. Comput.* 142 (2003) 1–16; MR1978243 (2004d:85001).
- [50] J.-H. He, Variational approach to the Lane-Emden equation, *Appl. Math. Comput.* 143 (2003) 539–541; MR1981715.
- [51] J. I. Ramos, Piecewise quasilinearization techniques for singular boundary-value problems, *Comput. Phys. Comm.* 158 (2004) 12–25; MR2105397 (2005f:65091).
- [52] J. I. Ramos, Linearization techniques for singular initial-value problems of ordinary differential equations, *Appl. Math. Comput.* 161 (2005) 525–542; MR2112422 (2005h:65101).
- [53] S. A. Yousefi, Legendre wavelets method for solving differential equations of Lane-Emden type, *Appl. Math. Comput.* 181 (2006) 1417–1422; MR2270772.
- [54] V. S. Ertürk, Differential transformation method for solving differential equations of Lane-Emden type, *Math. Comput. Appl.* 12 (2007) 135–139; MR2396843.
- [55] N. Guzel, M. Bayram and C. Guler, On the numerical solution of Lane-Emden differential equations with index- m , *Far East J. Appl. Math.* 29 (2007) 217–232; MR2372061.
- [56] A. Yildirim, A. and T. Özis, Solutions of singular IVPs of Lane-Emden type by homotopy perturbation method, *Phys. Lett. A* 369 (2007) 70–76.
- [57] C. M. Khalique and P. Ntsime, Exact solutions of the Lane-Emden-type equation, *New Astronomy* 13 (2008) 476–480.
- [58] H. R. Marzban, H. R. Tabrizidooz and M. Razzaghi, Hybrid functions for non-linear initial-value problems with applications to Lane-Emden type equations, *Phys. Lett. A* 372 (2008) 5883–5886.
- [59] J. I. Ramos, Series approach to the Lane-Emden equation and comparison with the homotopy perturbation method, *Chaos Solitons Fractals* 38 (2008) 400–408; MR2415942.
- [60] M. Dehghan and F. Shakeri, Approximate solution of a differential equation arising in astrophysics using the variational iteration method, *New Astronomy* 13 (2008) 53–59.

- [61] R. A. Van Gorder and K. Vajravelu, Analytic and numerical solutions to the Lane- Emden equation, *Phys. Lett. A* 372 (2008) 6060–6065.
- [62] M. S. H. Chowdhury and I. Hashim, Solutions of Emden-Fowler equations by homotopy-perturbation method, *Nonlinear Analysis: Real World Appl.* 10 (2009) 104–115.
- [63] S. Srivastava, A new solution of the Lane-Emden equation of index $n = 5$, *Astrophys. J.* 136 (1962) 680–681.
- [64] V. D. Sharma, A non-elliptic solution of the Lane-Emden equation for index $n = 5$, *Phys. Lett. A* 60 (1977) 381–382; MR0452446 (56 #10726).
- [65] J. O. Murphy, A finite radius solution for the polytrope index 5, *Proc. Astron. Soc. Austral.* 4 (1980) 37–39.
- [66] J. O. Murphy, On the F -type and M -type solutions of the Lane-Emden equation, *Proc. Astron. Soc. Austral.* 4 (1980) 41–43.
- [67] J. O. Murphy, A sequence of E -type composite analytical solutions of the Lane-Emden equation, *Proc. Astron. Soc. Austral.* 4 (1982) 376–378.
- [68] J. O. Murphy, Composite analytical solutions of the Lane-Emden equation with polytropic indices $n = 1$ and $n = 5$, *Proc. Astron. Soc. Austral.* 5 (1983) 175–179.
- [69] M. A. Lampert and R. U. Martinelli, Solution of the non-linear Poisson-Boltzmann equation in the interior of charged, spherical and cylindrical vesicles. I, The high-charge limit, *Chem. Phys.* 88 (1984) 399–413.
- [70] S. Chandrasekhar and G. W. Wares, The isothermal function, *Astrophys. J.* 109 (1949) 551–554.
- [71] F. K. Liu, Polytropic gas spheres: an approximate analytic solution of the Lane-Emden equation, *Monthly Notices Royal Astron. Soc.* 281 (1996) 1197–1205; arXiv:astro-ph/9512061.
- [72] P. Natarajan and D. Lynden-Bell, An analytic approximation to the isothermal sphere, *Monthly Notices Royal Astron. Soc.* 286 (1997) 268–270; arXiv:astro-ph/9604084v1.
- [73] A.-M. Wazwaz, A new algorithm for solving differential equations of Lane-Emden type, *Appl. Math. Comput.* 118 (2001) 287–310; MR1812960 (2001k:65107).

- [74] E. Momoniat and C. Harley, Approximate implicit solution of a Lane-Emden equation, *New Astronomy* 11(2006) 520–526.
- [75] N. Guzel and M. T. Oylumlu, Numerical solution of isothermal gas sphere problem, *Internat. J. Comput. Methods* 4 (2007) 163–171; MR2348108 (2008g:65095).