

Series-Parallel Networks

STEVEN FINCH

July 7, 2003

Series and parallel connections are usually first encountered in the study of electrical circuits. Our approach is to first examine a relevant class of partially ordered sets (posets) and then to define series-parallel networks by analogy [1]. Interesting asymptotic constants appear everywhere, similar to those associated with counting various species of trees [2]. We also talk briefly about the enumeration of Boolean (or switching) functions under different notions of equivalence.

0.1. Series-Parallel Posets. We introduce two procedures for combining two posets (S, \leq) and (S', \leq) to obtain a new poset, assuming that $S \cap S' = \emptyset$:

- the **disjoint sum** $S \oplus S'$ is the poset on $S \cup S'$ such that $x \leq y$ in $S \oplus S'$ if either $x, y \in S$ and $x \leq y$ in S , or $x, y \in S'$ and $x \leq y$ in S'
- the **linear product** $S \odot S'$ is the poset on $S \cup S'$ such that $x \leq y$ in $S \odot S'$ if $x, y \in S$ and $x \leq y$ in S , or $x, y \in S'$ and $x \leq y$ in S' , or $x \in S$ and $y \in S'$.

Clearly \oplus is commutative but \odot is not. A **series-parallel poset** is one that can be recursively constructed by applying the operations of disjoint sum and linear product, starting with a single point [3].

Define a poset to be **N-free** if there is no subset $\{a, b, c, d\}$ whose only nontrivial relations are given by

$$a < c, \quad a < d, \quad b < d.$$

It can be proved that a finite poset is series-parallel if and only if it is N-free [4, 5, 6]. Hence there are 15 series-parallel posets with 4 points (see the 16 posets in Figure 2 of [7] and eliminate the poset that looks like an “N”).

There are two cases we shall consider. The number f_n of unlabeled series-parallel posets with n points has (ordinary) generating function [3, 8, 9]

$$F(x) = \sum_{n=0}^{\infty} f_n x^n = 1 + x + 2x^2 + 5x^3 + 15x^4 + 48x^5 + 167x^6 + 602x^7 + 2256x^8 + \dots$$

⁰Copyright © 2003 by Steven R. Finch. All rights reserved.

which satisfies the functional equation

$$F(x) = \exp \left[\sum_{k=1}^{\infty} \frac{1}{k} \left(F(x^k) + \frac{1}{F(x^k)} + x^k - 2 \right) \right]$$

Alternatively, if the sequence $\{\hat{f}_n\}$ is defined by $1/F(x) = \sum_{n=0}^{\infty} \hat{f}_n x^n$, then

$$F(x) = \prod_{j=1}^{\infty} (1 - x^j)^{-(f_j + \hat{f}_j + \delta_{j,1})}$$

where $\delta_{j,k} = 1$ when $j = k$ and $\delta_{j,k} = 0$ otherwise. Using such properties, it follows that

$$f_n \sim \beta \cdot n^{-3/2} \cdot \alpha^{-n}$$

where $\alpha = 0.2163804273\dots$ is the unique positive root of $F(x) = \varphi$ and φ is the Golden mean, and where

$$\beta = \sqrt{\frac{1}{(3\sqrt{5} - 5)\pi} \left[\frac{\alpha}{1 - \alpha} + \sum_{i=2}^{\infty} \alpha^i F'(\alpha^i) \left(1 - \frac{1}{F(\alpha^i)^2} \right) \right]} = 0.2291846208\dots$$

The number g_n of labeled series-parallel posets with n points has (exponential) generating function [1, 3, 8, 9]

$$\begin{aligned} G(x) &= \sum_{n=1}^{\infty} \frac{g_n}{n!} x^n = x + \frac{3}{2!} x^2 + \frac{19}{3!} x^3 + \frac{195}{4!} x^4 + \frac{2791}{5!} x^5 + \frac{51303}{6!} x^6 + \frac{1152019}{7!} x^7 + \dots \\ &= \left(\ln(1+x) - \frac{x^2}{1+x} \right)^{\langle -1 \rangle} = \left(\sum_{k=1}^{\infty} (-1)^{k+1} \frac{k+1}{k} x^k \right)^{\langle -1 \rangle} \end{aligned}$$

where the notation $P(x)^{\langle -1 \rangle}$ denotes the reversion of the power series $P(x)$. Well-established theory [10, 11, 12] gives that

$$g_n \sim \eta \cdot n! \cdot n^{-3/2} \cdot \xi^{-n}$$

where $\xi = \ln(\varphi) - 2\varphi + 3 = 0.2451438475\dots$ and

$$\eta = \sqrt{\frac{\xi}{2\sqrt{5}(2 - \varphi)\pi}} = 0.2137301074\dots$$

Now let us define an equivalence relation on the set of series-parallel posets with n points, induced simply by declaring $S \odot S'$ and $S' \odot S$ to be equivalent. (See Figure 1.) The equivalence classes correspond to what are called **two-terminal series-parallel networks** with n edges [13, 14, 15, 16, 17, 18, 19], with the understanding that




Figure 1: There are 10 non-equivalent (unlabeled) series-parallel posets with 4 points. Note the analogy with Figure 2.

- points of a poset are mapped in a one-to-one manner to edges of the corresponding network
- two points of the poset are comparable if and only if the analogous edges of the network are connected in series
- two points of the poset are incomparable if and only if the analogous edges of the network are connected in parallel.

(See Figures 2 and 3.) The leftmost and rightmost points are the terminals (two distinguished points playing a role similar to that of the root of a rooted tree). A network, however, is not necessarily a graph since it may possess multiple parallel edges. Observe that an interchange of parts of the network, either in series or in parallel, is immaterial. In other words, when we count series-parallel networks, our tally is unaffected by a permutation of variables in the indicated Boolean representations.

0.2. Series-Parallel Networks. The number u_n of unlabeled series-parallel networks with n edges has generating function [20]

$$U(x) = \sum_{n=0}^{\infty} u_n x^n = 1 + x + 2x^2 + 4x^3 + 10x^4 + 24x^5 + 66x^6 + 180x^7 + 522x^8 + \dots$$

which satisfies the functional equation

$$U(x) = \exp \left[\sum_{k=1}^{\infty} \frac{1}{2k} (U(x^k) + x^k - 1) \right]$$




Figure 2: There are 10 unlabeled series-parallel networks with 4 edges, that is, $u_4 = 10$. The “essentially parallel” networks constitute the first row and the “essentially series” networks constitute the second row.




Figure 3: There are 8 labeled series-parallel networks with 3 edges, that is, $v_3 = 8$. The “essentially parallel” networks constitute the first row and the “essentially series” networks constitute the second row.

Alternatively, we have

$$U(x) = \prod_{j=1}^{\infty} (1 - x^j)^{-(u_j + \delta_{j,1})/2}$$

Using these properties, it follows that [15, 21, 22, 23]

$$u_n \sim \lambda \cdot n^{-3/2} \cdot \kappa^{-n}$$

where $\kappa = 0.2808326669\dots = (3.5608393095\dots)^{-1}$ is the unique positive root of $U(x) = 2$ and

$$\lambda = \sqrt{\frac{1}{\pi} \left[\frac{\kappa}{1 - \kappa} + \sum_{i=2}^{\infty} \kappa^i U'(\kappa^i) \right]} = 0.4127628892\dots = 2 \cdot (0.2063814446\dots)$$

This also gives the number of non-equivalent Boolean functions of n variables, built only with $+$ (disjunction) and \cdot (conjunction).

The number v_n of labeled series-parallel networks with n edges has generating function [1, 24]

$$\begin{aligned} V(x) &= \sum_{n=1}^{\infty} \frac{v_n}{n!} x^n = x + \frac{2}{2!} x^2 + \frac{8}{3!} x^3 + \frac{52}{4!} x^4 + \frac{472}{5!} x^5 + \frac{5504}{6!} x^6 + \frac{78416}{7!} x^7 + \dots \\ &= (2 \ln(1+x) - x)^{(-1)} = \left(\sum_{k=1}^{\infty} (-1)^{k+1} \frac{2}{k} x^k \right)^{(-1)} \end{aligned}$$

By techniques similar to those used to analyze $\{g_n\}$, we have [21, 25]

$$v_n \sim \tau \cdot n! \cdot n^{-3/2} \cdot \sigma^{-n}$$

where $\sigma = 2 \ln(2) - 1 = 0.3862943611\dots = (2.5886994495\dots)^{-1}$ and

$$\tau = \sqrt{\frac{\sigma}{\pi}} = 0.3506584008\dots = 2 \cdot (0.1753292004\dots)$$

Related work involves bracketing of n -symbol products [26] and phylogenetic trees [27].

0.3. Series-Parallel Networks Without Multiple Parallel Edges. If we prohibit multiple parallel edges, so that the networks under consideration are all graphs, different constants arise. (See Figure 4).

The number q_n of such unlabeled series-parallel networks with n edges has generating function [28]

$$Q(x) = \sum_{n=0}^{\infty} q_n x^n = 1 + x + x^2 + 2x^3 + 4x^4 + 8x^5 + 18x^6 + 40x^7 + 94x^8 + 224x^9 + \dots$$




Figure 4: There are 8 unlabeled series-parallel networks with 5 edges that obey the prohibition against multiple parallel edges, that is, $q_5 = 8$. The “essentially parallel” networks constitute the first row and the “essentially series” networks constitute the second row.

which satisfies the functional equation

$$Q(x) = \exp \left[\sum_{k=1}^{\infty} \frac{1}{2k} \left(Q(x^k) - x^{2k} + x^k - 1 \right) \right]$$

Alternatively, we have

$$Q(x) = \prod_{j=1}^{\infty} \left(1 - x^j \right)^{-(q_j + \delta_{j,1} - \delta_{j,2})/2}$$

Using these properties, it follows that [21]

$$q_n \sim \nu \cdot n^{-3/2} \cdot \mu^{-n}$$

where $\mu = 0.3462834070\dots$ is the unique positive root of $Q(x) = 2$ and

$$\nu = \sqrt{\frac{1}{\pi} \left[\frac{\mu}{1 + \mu} + \sum_{i=2}^{\infty} \mu^i Q'(\mu^i) \right]} = 0.3945042461\dots = 2 \cdot (0.1972521230\dots)$$

The number r_n of such labeled series-parallel networks with n edges has generating function [29]

$$\begin{aligned} R(x) &= \sum_{n=1}^{\infty} \frac{r_n}{n!} x^n = x + \frac{1}{2!} x^2 + \frac{4}{3!} x^3 + \frac{20}{4!} x^4 + \frac{156}{5!} x^5 + \frac{1472}{6!} x^6 + \frac{17396}{7!} x^7 + \dots \\ &= \left((x + 1)^2 \exp(-x) - 1 \right)^{\langle -1 \rangle} = \left(\sum_{k=1}^{\infty} (-1)^k \frac{k^2 - 3k + 1}{k!} x^k \right)^{\langle -1 \rangle} \end{aligned}$$

Proceeding as before, we have [21]

$$r_n \sim \omega \cdot n! \cdot n^{-3/2} \cdot \theta^{-n}$$

where $\theta = 4/e - 1 = 0.4715177646\dots$ and

$$\omega = \frac{1}{2} \sqrt{\frac{e\theta}{\pi}} = 0.3193679560\dots = 2 \cdot (0.1596839780\dots)$$

It follows that the probability that a random n -edge series-parallel network has no multiple parallel edges is asymptotically

$$\left(\frac{\nu}{\lambda}\right) \left(\frac{\kappa}{\mu}\right)^n = (0.9557648142\dots)(0.8109908278\dots)^n$$

if the network is unlabeled and

$$\left(\frac{\omega}{\tau}\right) \left(\frac{\sigma}{\theta}\right)^n = (0.9107665899\dots)(0.8192572794\dots)^n$$

if the network is labeled. We hope to report on later on other relevant material in [21].

0.4. Boolean Functions. We have already enumerated the number u_n of distinct Boolean functions of n variables, built only with $+$ and \cdot , under the action of the symmetric group S_n .

Of course, the set of *all* Boolean functions also includes those involving complementation of variables ($\neg X$). Let us examine briefly this larger set [30, 31]. Define two Boolean functions to be **equivalent** if they are identical up to a bijective renaming of the variables. The number of equivalence classes in this case is asymptotically [32, 33, 34]

$$2^{2^n} / n!$$

hence no new constants arise. Define two Boolean functions to be **congruent** if they are identical up to a bijective renaming of the variables and an additional complementation of some of the variables. The number of congruence classes is asymptotically

$$2^{2^n - n} / n!$$

Other results of this kind are also known, but none contain new constants.

Let us return to our original set of Boolean functions of n variables and let \mathbb{F}_2 denote the binary field. S_n is a subgroup of the group T_n of invertible linear transformations $\mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$, namely, the $n \times n$ matrices that have exactly one 1 in each row and each column. What can be said about the number \tilde{u}_n of distinct Boolean functions, built only with $+$ and \cdot , under the action of the (larger) group T_n ? Our experience with u_n leads us to conjecture that the asymptotics of \tilde{u}_n will be quite interesting.

0.5. Irreducible Posets. Another unsolved problem involves the number a_n of unlabeled (\oplus, \odot) -irreducible posets with n points. It is known that

$$A(x) = \sum_{n=0}^{\infty} a_n x^n = x + x^4 + 12x^5 + 104x^6 + 956x^7 + 10037x^8 + 126578x^9 + 1971005x^{10} + \dots$$

and, further, that

$$P(x) = \exp \left[\sum_{k=1}^{\infty} \frac{1}{k} \left(P(x^k) + \frac{1}{P(x^k)} + A(x^k) - 2 \right) \right]$$

where

$$P(x) = \sum_{n=0}^{\infty} p_n x^n = 1 + x + 2x^2 + 5x^3 + 16x^4 + 63x^5 + 318x^6 + 2045x^7 + 16999x^8 + \dots$$

is the generating function of (arbitrary) unlabeled posets [3, 7, 9]. What can be said about the asymptotics of a_n ? Even a nice functional equation for $A(x)$ in-and-by-itself is probably impossible.

Addendum. Bodirsky, Giménez, Kang & Noy [35] have determined the asymptotic number of labelled series-parallel graphs, which we hope to report on later.

REFERENCES

- [1] R. P. Stanley, *Enumerative Combinatorics*, v 2., Cambridge Univ. Press, 1999, pp. 88–89, 133; MR1676282 (2000k:05026).
- [2] S. R. Finch, Otter’s tree enumeration constants, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 295–316.
- [3] R. P. Stanley, Enumeration of posets generated by disjoint unions and ordinal sums, *Proc. Amer. Math. Soc.* 45 (1974) 295–299; MR0351928 (50 #4416).
- [4] J. Valdes, R. E. Tarjan, and E. L. Lawler, The recognition of series parallel digraphs, *Proc. 11th ACM Symp. on Theory of Computing (STOC)*, Atlanta, ACM, 1979, pp. 1–12; *SIAM J. Comput.* 11 (1982) 298–313; MR0564616 (81d:68088) and MR0652904 (84d:68073)
- [5] J. Grabowski, On partial languages, *Fundamenta Informaticae* 4 (1981) 427–498; MR0645249 (83i:68107).
- [6] J. L. Gischer, The equational theory of pomsets, *Theoret. Comput. Sci.* 61 (1988) 199–224; MR0980242 (90b:68058).

- [7] S. R. Finch, Transitive relations, topologies and partial orders, unpublished note (2003).
- [8] M. H. El-Zahar, Enumeration of ordered sets, *Algorithms and Order*, Proc. NATO Advanced Study Institute, Ottawa, 1987, ed. I. Rival, Kluwer, 1989, pp. 327–352; MR1037788 (90m:06005).
- [9] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A003430, A003431, A046906, A053554.
- [10] E. A. Bender, Asymptotic methods in enumeration, *SIAM Rev.* 16 (1974) 485-515; errata 18 (1976) 292; MR0376369 (51 #12545) and MR0437344 (55 #10276).
- [11] A. Meir and J. W. Moon, On an asymptotic method in enumeration, *J. Combin. Theory Ser. A* 51 (1989) 77–89; erratum 52 (1989) 163; MR0993650 (90e:05007a) and MR1008169 (90e:05007b).
- [12] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, unpublished manuscript, 2001.
- [13] P. A. MacMahon, Yoke-chains and multipartite compositions in connexion with the analytical forms called “trees”, *Proc. London Math. Soc.* 22 (1891) 330-346; *Collected Papers*, v. 1, ed. G. E. Andrews, MIT Press, 1978, pp. 600-616.
- [14] P. A. MacMahon, The combinations of resistances, *The Electrician* 28 (1892) 601-602; *Collected Papers*, v. 1, ed. G. E. Andrews, MIT Press, 1978, pp. 617-619.
- [15] J. Riordan and C. E. Shannon, The number of two-terminal series-parallel networks, *J. Math. Phys. Mass. Inst. Tech.* 21 (1942) 83–93; *Claude Elwood Shannon: Collected Papers*, ed. N. J. A. Sloane and A. D. Wyner, IEEE Press, 1993, pp. 560-570; MR0007511 (4,151c).
- [16] W. Knödel, Über Zerfällungen, *Monatsh. Math.* 55 (1951) 20–27; MR0040996 (12,778d).
- [17] L. Carlitz and J. Riordan, The number of labeled two-terminal series-parallel networks, *Duke Math. J.* 23 (1956) 435–445; MR0078940 (18,3e).
- [18] J. Riordan, *An Introduction to Combinatorial Analysis*, Wiley, 1958, pp. 139-143; MR0096594 (20 #3077).

- [19] Z. A. Lomnicki, Two-terminal series-parallel networks, *Adv. Appl. Probab.* 4 (1972) 109–150; MR0319662 (47 #8204).
- [20] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A000084, A000669, A058964, A058965.
- [21] J. W. Moon, Some enumerative results on series-parallel networks, *Random Graphs '85*, Proc. 1985 Poznan conf, ed. M. Karonski and Z. Palka, Annals of Discrete Math. 33, North-Holland, 1987, pp. 199–226; MR0930492 (89h:05004).
- [22] O. Golinelli, Asymptotic behavior of two-terminal series-parallel networks, [cond-mat/9707023](https://arxiv.org/abs/cond-mat/9707023).
- [23] E. M. Rains and N. J. A. Sloane, The series-parallel constant, unpublished note (2001).
- [24] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A000311, A006351.
- [25] A. Meir and J. W. Moon, The asymptotic behaviour of coefficients of powers of certain generating functions, *European J. Combin.* 11 (1990) 581–587; MR1078714 (91m:05014).
- [26] J. Riordan, The blossoming of Schröder's fourth problem, *Acta Math.* 137 (1976) 1–16; MR0429582 (55 #2593).
- [27] L. R. Foulds and R. W. Robinson, Enumeration of phylogenetic trees without points of degree two, *Ars Combin.* 17A (1984) 169–183; MR0746182 (85f:05045).
- [28] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A058385, A058386, A058387.
- [29] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A058379, A058380, A058381.
- [30] B. Borchert, D. Ranjan, and F. Stephan, On the computational complexity of some classical equivalence relations on Boolean functions, *Theory Comput. Syst.* 31 (1998) 679–693; available at <http://www.ubka.uni-karlsruhe.de/cgi-bin/psview?document=/1995/informatik/24>; MR1642457 (2000i:68055).
- [31] P. Savický and A. R. Woods, The number of Boolean functions computed by formulas of a given size, *Random Structures Algorithms* 13 (1998) 349–382; MR1662790 (2000e:68077).

- [32] M. A. Harrison, On asymptotic estimates in switching and automata theory, *J. ACM* 13 (1966) 151–157; MR0186504 (32 #3964).
- [33] M. A. Harrison, Counting theorems and their applications to classification of switching functions, *Recent Developments in Switching Theory*, ed. A. Mukhopadhyay, Academic Press, 1971, pp. 85–120; MR0280279 (43 #6000).
- [34] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A000231, A003180, A000616.
- [35] M. Bodirsky, O. Giménez, M. Kang and M. Noy, Enumeration and limit laws of series-parallel graphs, math.CO/0512435.