

Planar Harmonic Mappings

STEVEN FINCH

March 29, 2005

Let D denote the open unit disk. A function $f : D \rightarrow \mathbb{C}$ is **planar harmonic** if it can be written as $f(z) = u(x, y) + i v(x, y)$ where $z = x + i y$ and where $u : D \rightarrow \mathbb{R}$, $v : D \rightarrow \mathbb{R}$ are harmonic, that is, are twice continuously differentiable and obey

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0, \quad \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = 0.$$

It can be shown that f is planar harmonic if and only if $f = g + \bar{h}$, where g, h are analytic on D and the overbar indicates complex conjugation ($\bar{z} = x - i y$).

Of course, a planar harmonic function f is analytic if and only if u and v are harmonic conjugates, that is, the Cauchy-Riemann equations

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

are satisfied. We are interested, in this essay, in functions f whose real and imaginary parts are not necessarily conjugate [1].

It turns out that f may be written as a twice continuously differentiable function of z and \bar{z} ; we abuse notation and use the same letter f to represent the new function. The Cauchy-Riemann equations become a single concise equation:

$$\frac{\partial f}{\partial \bar{z}} = 0$$

and the condition that Laplacians vanish becomes

$$4 \frac{\partial^2 f}{\partial z \partial \bar{z}} = 0.$$

Thus the expression f is independent of \bar{z} for analytic functions f , and the expression $\partial f / \partial z$ is independent of \bar{z} for planar harmonic functions f .

A planar harmonic function $f : D \rightarrow \mathbb{C}$ is a **mapping** if it is one-to-one. Hence the class of planar harmonic mappings includes the subclass of univalent functions we have studied elsewhere [2, 3, 4, 5]. Define also the **dilatation** of f

$$\omega = \frac{\overline{\partial f}}{\partial \bar{z}} / \frac{\partial f}{\partial z}$$

which will be needed later.

⁰Copyright © 2005 by Steven R. Finch. All rights reserved.

0.1. Heinz’s Inequality. We consider here planar harmonic mappings f that map D onto D , with the property that $f(0) = 0$. Heinz [6] proved that

$$\left| \frac{\partial f}{\partial z}(0, 0) \right|^2 + \left| \frac{\partial f}{\partial \bar{z}}(0, 0) \right|^2 \geq c$$

for some constant $c \geq 0.1788 = 0.3576/2$. The lower bound was improved to $0.32 = 0.64/2$ by Nitsche [7, 8], $0.4345 = 0.8691/2$ by de Vries [9], $0.4476 = 0.8952/2$ by Nitsche [10], $0.6411 = 1.2822/2$ by de Vries [11], and $0.6584 = 1.3168/2$ by Wegmann [12]. The conjecture that

$$c = \frac{27}{4\pi^2} = 0.6839179895\dots = \frac{1}{2}(1.3678359791\dots)$$

mentioned by Wegmann [12] seems to have been anticipated by Hopf [13]. A proof of this conjecture was first given by Hall [1, 14]; the extremal function is achieved via approximations $D \rightarrow D$ of a mapping $D \rightarrow T$, where T is an inscribed equilateral triangle, with dilatation $\omega(z) = z$.

Hall’s proof involves the Fourier coefficients of homeomorphisms $C \rightarrow C$ of the unit circle C . Some related problems are given in [14]; one of these has been solved [15]. Heinz [16] also proved the inequality

$$\left| \frac{\partial f}{\partial z}(z, \bar{z}) \right|^2 + \left| \frac{\partial f}{\partial \bar{z}}(z, \bar{z}) \right|^2 \geq \frac{1}{\pi^2}$$

which is valid for all $z \in D$; improvements in special cases appear in [17, 18].

0.2. Minimal Surfaces. Consider a minimal surface over the unit disk D of the form

$$\{(x, y, z) \in \mathbb{R}^3 : z = F(x, y), (x, y) \in D\}$$

and let κ denote its Gaussian curvature at the origin. In words, the surface is locally area-minimizing: Each suitable small piece of it has the least possible area for any surface spanning the boundary of that piece. By the calculus of variations, we have the nonlinear PDE

$$\left[1 + \left(\frac{\partial F}{\partial y} \right)^2 \right] \frac{\partial^2 F}{\partial x^2} - 2 \frac{\partial F}{\partial x} \frac{\partial F}{\partial y} \frac{\partial^2 F}{\partial x \partial y} + \left[1 + \left(\frac{\partial F}{\partial x} \right)^2 \right] \frac{\partial^2 F}{\partial y^2} = 0;$$

hence the mean curvature of the surface is everywhere zero. A precise determination of F is difficult – this is called **Plateau’s problem** – but nature solves it effortlessly, as can be demonstrated by dipping a bent wire loop in a soap solution [19, 20].

A consequence of Heinz's inequality [6] is that

$$|\kappa| \leq \frac{4}{c} = \frac{16\pi^2}{27} = 5.8486544599\dots$$

by Hall's theorem [14], but this is not sharp. In fact, it is conjectured that [1]

$$|\kappa| \leq \frac{\pi^2}{2} = 4.9348022005;$$

this has however been proved only in the special case that the minimal surface has a horizontal tangent plane at the origin [21]. A general proof could be obtained utilizing the following.

Consider planar harmonic mappings f that map D onto D , with the two properties that $f(0) = 0$ and ω is the square of an analytic function. (Note that this final requirement is not met by $\omega(z) = z$.) Hall [22] recently computed that

$$\left| \frac{\partial f}{\partial z}(0,0) \right|^2 + \left| \frac{\partial f}{\partial \bar{z}}(0,0) \right|^2 \geq \tilde{c}$$

for some constant $\tilde{c} > c + 10^{-5}/2$. It is conjectured that $\tilde{c} = 8/\pi^2$ (from which $4/\tilde{c} = \pi^2/2$ would proceed immediately). The expected extremal function is a mapping $D \rightarrow S$, where S is an inscribed square, with dilatation $\omega(z) = z^2$. A proof that $\tilde{c} = 8/\pi^2$ would be a major step forward in understanding minimal surfaces. See [23] for more open questions.

0.3. Soap Films. As an aside, we give an elementary problem [24, 25]. Consider the catenoid-shaped soap film formed between two parallel rings centered at $(-\xi, 0, 0)$ and $(\xi, 0, 0)$ and of unit radius, where $\xi > 0$ is suitably small. If the rings are slowly pulled apart (that is, if ξ increases), there is a certain threshold at which the minimal surface becomes unstable and is likely to collapse to a disjoint union of two disks. More precisely, if $\xi < \xi_0 = 0.5276973969\dots$, then the catenoid corresponds to the global minimum for surface area while the two-disk configuration corresponds to only a local minimum. Here ξ_0 and $a = 0.8255174536\dots$ are solutions of the simultaneous equations

$$\begin{cases} a \cosh\left(\frac{\xi_0}{a}\right) = 1, \\ 2\pi a^2 \sinh\left(\frac{\xi_0}{a}\right) \cosh\left(\frac{\xi_0}{a}\right) + 2\pi a \xi_0 = 2\pi. \end{cases}$$

If $\xi > \xi_0$, then the two-disk configuration corresponds to the global minimum while the catenoid corresponds to only a local minimum for $\xi < \xi_1 = 0.6627434193\dots$; no

such catenoid exists for $\xi > \xi_1$. Here ξ_1 and $b = 0.5524341245\dots$ are solutions of the simultaneous equations

$$\begin{cases} b \cosh\left(\frac{\xi_1}{b}\right) = 1, \\ \cosh\left(\frac{\xi_1}{b}\right) - \frac{\xi_1}{b} \sinh\left(\frac{\xi_1}{b}\right) = 0. \end{cases}$$

Interestingly, we have seen the value for ξ_1 before: In [26], it arose in a different context altogether and was called the *Laplace limit constant*.

REFERENCES

- [1] P. Duren, *Harmonic Mappings in the Plane*, Cambridge Univ. Press, 2004, pp. 1–7, 18–23, 66–72, 182–186; MR2048384 (2005d:31001).
- [2] S. R. Finch, Bloch-Landau constants, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 456–459.
- [3] S. R. Finch, Littlewood-Clunie-Pommerenke constants, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 470–473.
- [4] S. R. Finch, Radii in geometric function theory, unpublished note (2004).
- [5] S. R. Finch, Coefficient estimates for univalent functions, unpublished note (2004).
- [6] E. Heinz, Über die Lösungen der Minimalflächengleichung, *Nachr. Akad. Wiss. Göttingen. Math.-Phys. Kl. Math.-Phys.-Chem. Abt.* (1952) 51–56; MR0054182 (14,885c).
- [7] J. C. C. Nitsche, On harmonic mappings, *Proc. Amer. Math. Soc.* 9 (1958) 268–271; MR0104233 (21 #2990a).
- [8] J. C. C. Nitsche, On the constant of E. Heinz, *Rend. Circ. Mat. Palermo* 8 (1959) 178–181; MR0113054 (22 #3895).
- [9] H. L. de Vries, A remark concerning a lemma of Heinz on harmonic mappings, *J. Math. Mech.* 11 (1962) 469–471; MR0142769 (26 #338).
- [10] J. C. C. Nitsche, Zum Heinzschen Lemma über harmonische Abbildungen, *Arch. Math.* 14 (1963) 407–410; MR0160919 (28 #4128).
- [11] H. L. de Vries, Über Koeffizientenprobleme bei Eiliniern und über die Heinzsche Konstante, *Math. Z.* 112 (1969) 101–106; MR0254742 (40 #7949).

- [12] R. Wegmann, Extremalfiguren für eine Klasse von isoperimetrischen Problemen, *Math. Nachr.* 69 (1975) 173–190; MR0428198 (55 #1223).
- [13] E. Hopf, On an inequality for minimal surfaces $z = z(x, y)$, *J. Rational Mech. Anal.* 2 (1953) 519–522; correction 2 (1953) 801–802; MR0055735 (14,1119d) and MR0058268 (15,348a).
- [14] R. R. Hall, On an inequality of E. Heinz, *J. d'Analyse Math.* 42 (1982/83) 185–198; MR0729409 (85k:42011).
- [15] A. Weitsman, On the Fourier coefficients of homeomorphisms of the circle, *Math. Res. Lett.* 5 (1998) 383–390; MR1637852 (99h:31003).
- [16] E. Heinz, On one-to-one harmonic mappings, *Pacific J. Math.* 9 (1959) 101–105; MR0104933 (21 #3683).
- [17] D. Partyka and K. Sakan, On Heinz's inequality, *Bull. Soc. Sci. Lett. Lodz Sér. Rech. Déform.* 37 (2002) 27–34; available online at http://math01.sci.osaka-cu.ac.jp/preprint/index_e.html; MR2055781 (2005e:31003).
- [18] D. Partyka and K. Sakan, On an asymptotically sharp variant of Heinz's inequality, unpublished note (2004).
- [19] J. C. C. Nitsche, *Vorlesungen über Minimalflächen*, Springer-Verlag, 1975; revised Engl. transl., *Lectures on Minimal Surfaces*, v. I, Cambridge Univ. Press, 1989; MR0448224 (56 #6533) and MR1015936 (90m:49031).
- [20] U. Dierkes, S. Hildebrandt, A. Küster and O. Wohlrab, *Minimal Surfaces*, v. I and II, Springer-Verlag, 1992; MR1215267 (94c:49001a) and MR1215268 (94c:49001b).
- [21] R. Finn and R. Osserman, On the Gauss curvature of non-parametric minimal surfaces, *J. d'Analyse Math.* 12 (1964) 351–364; MR0166694 (29 #3967).
- [22] R. R. Hall, The Gaussian curvature of minimal surfaces and Heinz' constant, *J. Reine Angew. Math.* 502 (1998) 19–28; MR1647626 (99j:53009).
- [23] J. Clunie and T. Sheil-Small, Harmonic univalent functions, *Annales Acad. Sci. Fenn. Ser. A I Math.* 9 (1984) 3–25; MR0752388 (85i:30014).
- [24] G. A. Bliss, *Calculus of Variations*, Math. Assoc. Amer., 1925, pp. 5–8, 85–127.
- [25] J. Oprea, *The Mathematics of Soap Films: Explorations with Maple*® , Amer. Math. Soc., 2000, pp. 11–12, 71–72, 181–191; MR1779498 (2002c:53013).

- [26] S. R. Finch, Laplace limit constant, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 266–268.