

Impatiemment Attendu

Robert Sedgewick
Princeton University

A
CONFERENCE
ON
THEORETICAL COMPUTER SCIENCE



August 15 – 17, 1977
University of Waterloo
Waterloo, Ontario
Canada

August 15 – 17, 1977

TABLE OF CONTENTS (continued)

Session III - continued

	<u>Page</u>
A New Complexity Measure for Languages by Luc Boasson, Bruno Courcelle and Maurice Nivat	130
Complexity of Grammatical Similarity Relations by H.B. Hunt, III and D.J. Rosenkrantz	139

Session IV: A. Borodin, Chairman

Addition Chain Methods for the Evaluation of Specific Polynomials by David Dobkin and Richard Lipton	146
Fast Algorithms for Composition and Reversion of Multivariate Power Series by R.P. Brent and H.T. Kung	149
Average Lower Bounds for Open-Addressing Hash Coding by Gaston H. Gonnet	159
Data Movement in Odd-Even Merging by Robert Sedgwick	163
A Fast Selection Algorithm and the Problem of Optimum Distribution of Effort by Zvi Galil and Nimrod Megiddo	173

Wednesday, August 17

Session V: Z. Manna, Chairman

Orthogonal Information Structures by Armin Cremers and Thomas N. Hibbard	182
Relational Equations, Grammars, and Programs by M.H. van Emden	191
Verification Decidability of Presburger Array Programs by Norihisa Suzuki and David Jefferson	202
Demi-Groupes de Matrices Localement Testables Characterisation et Decidabilite by G. Jacob	213
Straight-Line Programs to Compute Finite Languages by Glenn B. Goodrich, Richard E. Ladner and Michael J. Fischer	221
A Graphical Proof Procedure for First-Order Logic by Philip T. Cox	230

Session VI: P.C. Fischer, Chairman

Automata with Data Storage by Jonathan Goldstine	239
Codes Asynchrones by Dominique Perrin	247
Toward a Theory of Data Encoding by Arnold L. Rosenberg and Lawrence Snyder	255
Aspects of the Database Security Problem by David Dobkin, Richard J. Lipton and Steven P. Reiss	262
Some Computational Problems Related to Database Concurrency Control by Christos H. Papadimitriou, Philip A. Bernstein and James B. Rothnie	275
AUTHOR INDEX	283

**Data Movement in Odd-Even Merging
by Robert Sedgwick**

$$\zeta(2z, a) = \frac{1}{2} - a + z(2 \ln \Gamma(a) - \ln 2\pi) + O(z^2).$$

The expansion for $\Gamma(z)$ is well known (see [1, eq. 6.1.33]), the next two expansions are elementary, and the expansion for the ζ -function follows directly from [16, p. 271]. Multiplying these series together, we find the expansion at $z=0$ of the integrand in (23):

$$\frac{1}{4z^2 \ln 2} + \frac{1}{4} \lg j - \frac{\gamma}{4 \ln 2} - \frac{1}{4} + \lg \frac{\Gamma(1/4)^2}{2\pi} + O(1). \quad (24)$$

This gives the residue at $z=0$ (the coefficient of $1/z$).

To this we must add the residue at the other poles of $1/(4z-1)$. The effect of these other terms is small (but not insignificant), and we shall encapsulate them in a single term,

$$\epsilon(j) \equiv \frac{2}{\ln 4} \sum_{k \neq 0} \Gamma\left(\frac{2k\pi i}{\ln 4}\right) j^{2k\pi i / \ln 4} \zeta\left(\frac{4k-1}{4}\right) = \sum_{k=1}^{\infty} (\xi_k \cos(k\pi \lg j) - \eta_k \sin(k\pi \lg j)),$$

where

$$\frac{2}{\ln 2} \Gamma\left(\frac{k\pi i}{\ln 2}\right) \zeta\left(\frac{2k\pi i}{\ln 2} + \frac{1}{4}\right) \equiv \xi_k + i\eta_k.$$

To finish the evaluation of our b_j and B_N we need to evaluate the Γ and ζ functions at these points along the imaginary axis (see [6, secs. 6.3 and 6.4]). Table 3 shows the values of ξ_k and η_k for $k=1, 2, 3$ computed in this way. The values get exceedingly small for larger k , as can be verified from the bounds (21) and (22).

Adding all the residues, we have, from (23):

$$b_j = \frac{1}{4} \lg j + \lg \frac{\Gamma(1/4)^2}{2\pi} - \frac{1}{4} - \frac{\gamma}{4 \ln 2} + \epsilon(j) + O(j^{-1/2}). \quad (25)$$

This leads to our final result.

Theorem 3. The average number of exchanges used by Batchier's odd-even merge for a random 2-ordered file of $2N$ elements is

$$B_N = \frac{1}{4} N \lg N + \left(\lg \frac{\Gamma(1/4)^2}{2\pi} + \frac{1}{4} - \frac{\gamma+2}{4 \ln 2} + \delta(N)\right) N + O(\sqrt{N} \log N),$$

where $\delta(N)$ is an oscillatory function which is

k	ξ_k	η_k
1	.003704670+	.002500177+
2	.000001560+	-.000000832-
3	.000000001-	.000000002+

$$\xi_k + i\eta_k = \frac{2}{\ln 2} \Gamma\left(\frac{k\pi i}{\ln 2}\right) \zeta\left(\frac{k\pi i}{\ln 2} + \frac{1}{4}\right)$$

$$\begin{aligned} \Gamma\left(\frac{1}{4}\right) &= 3.6256099082+ \\ \frac{1}{\ln 2} &= 1.4426950408+ \\ \gamma &= 0.5772156649+ \\ \pi &= 3.1415926535+ \end{aligned}$$

Table 3. Values of constants

constant if N is a power of two, with $|\delta(N)| < .000680$, and $\delta(2^n) \approx .000317(-1)^n$. (The constant

$$\alpha \equiv \lg \frac{\Gamma(1/4)^2}{2\pi} + \frac{1}{4} - \frac{\gamma+2}{4 \ln 2}$$

has the approximate value .385417224.)

Proof. From the discussion above, we need only substitute our result (25) for b_j into our eq. (14) for B_N and perform the summation. We have

$$B_N = \frac{1}{2} \frac{4^N}{\binom{2N}{N}} \sum_{1 \leq j < N} \frac{\binom{2j}{j}}{4^j} \left(\frac{1}{4} \lg j + \alpha + \frac{1}{2 \ln 2} + \epsilon(j) + O(j^{-1/2})\right).$$

The $\alpha + 1/(2 \ln 2)$ terms are easily taken care of. Working backwards from (14), we see that they contribute $(\alpha + 1/(2 \ln 2))N$ to B_N . (The sum may be evaluated directly as an interesting application of several identities in Knuth [13]; a simple induction could also be used.)

For the other terms, we can remove the binomial coefficients with Stirling's approximation, as in the derivation of (16). We have

$$\frac{\binom{2j}{j}}{4^j} = \frac{1}{\sqrt{\pi j}} + O(j^{-3/2}), \quad \frac{4^N}{\binom{2N}{N}} = \sqrt{\pi N} + O(N^{-1/2}).$$

Therefore the $O(j^{-1/2})$ term sums to $O(\sqrt{N} \log N)$, and

$$\begin{aligned} \frac{1}{2} \frac{4^N}{\binom{2N}{N}} \sum_{1 \leq j < N} \frac{\binom{2j}{j}}{4^j} \frac{1}{4} \lg j &= \frac{\sqrt{N}}{8} \sum_{1 \leq j < N} \frac{\lg j}{\sqrt{j}} + O(\sqrt{N}) \\ &= \frac{\sqrt{N}}{8} \int_1^N \frac{\lg x}{\sqrt{x}} dx + O(\sqrt{N}) \\ &= \frac{1}{4} N \lg N - \frac{1}{2 \ln 2} N + O(\sqrt{N}). \end{aligned}$$

Here the second step follows from Euler-McLaurin summation (see, for example, [14, p. 110]) and the integral is evaluated by making the substitution $x = y^2$.

It remains to evaluate the oscillatory term

$$N\delta(N) = \frac{1}{2} \frac{4^N}{\binom{2N}{N}} \sum_{1 \leq j < N} \frac{\binom{2j}{j}}{4^j} \epsilon(j).$$

After substituting for $\epsilon(j)$, we proceed in the same way as we did for the $\lg j$ term. The result of using Stirling's approximation on the binomial coefficients and Euler-McLaurin summation on the resulting sums is

$$\begin{aligned} \delta(N) &= \frac{1}{2\sqrt{N}} \sum_{k \geq 1} (\xi_k \int_1^N \frac{\cos k\pi \lg x}{\sqrt{x}} dx \\ &\quad - \eta_k \int_1^N \frac{\sin k\pi \lg x}{\sqrt{x}} dx) + O(\sqrt{N}). \end{aligned}$$

These integrals are elementary; the substitutions $x = y^2$, then $t = 2\pi k \lg y$ transform them into standard integrals (for example, [1, Eqs. 4.3.136, 4.3.137]) with the eventual result


$$\begin{aligned} \delta(N) &= \sum_{k \geq 1} \frac{\sigma_k}{2\sigma_k + 1} (\xi_k (\sigma_k \cos \pi k \lg N + \sin \pi k \lg N) \\ &\quad - \eta_k (\sigma_k \sin \pi k \lg N - \cos \pi k \lg N)) \end{aligned}$$

where σ_k is $(\ln 2/2\pi k)$. From this formula, we see that $\delta(N)$ is constant when N is a power of two, and has an oscillatory nature between powers of two. With the aid of Table 3 we can easily compute the stated values. \square

$$\frac{2}{\ln 2} \Gamma\left(\frac{k\pi i}{\ln 2}\right) \zeta\left(\frac{2k\pi i}{\ln 2} + \frac{1}{4}\right)$$

18th Annual
Symposium on Foundations
of Computer Science

(Formerly called the Annual Symposium on Switching and Automata Theory)




OCTOBER 31-NOVEMBER 2, 1977

OCTOBER 31-NOVEMBER 2, 1977

IEEE 77 CH1278-1 C

 IEEE COMPUTER SOCIETY

 INSTITUTE OF ELECTRICAL AND ELECTRONICS ENGINEERS

Session III - continued

On Uniquely Representable Data Structures
by Lawrence Snyder 142

Finite Automata in a 2-Dimensional Space
by Manuel Blum and William Sakoda 147

Session IV: Sam Winograd, Chairman

Applications of a Planar Separator Theorem
by Richard J. Lipton and Robert E. Tarjan 162

The Power of Commutativity
by L. Hyafil 171

On Taking Roots in Finite Fields
by Leonard Adleman, Kenneth Manders, and Gary Miller 175

Saving Space in Fast String-Matching
by Zvi Galil and Joel Seiferas 179

A New Proof of the Linearity of the Boyer-Moore String Searching Algorithm
by Leo J. Guibas and Andrew M. Odlyzko 189

On the Average Number of Registers Required for Evaluating Arithmetic Expressions
by P. Flajolet, J. C. Raoult, and J. Vuillemin 196

Wednesday, November 2

Session V: Nancy Lynch, Chairman

Fast Approximation Algorithms for Knapsack Problems
by E. L. Lawler 206

Combinatorial Analysis of an Efficient Algorithm for Processor and Storage Allocation
by E. G. Coffman and Joseph Y-T. Leung 214

Probabilistic Computations - Toward a Unified Measure of Complexity
by Andrew C. Yao 222

On Triangulations of a Set of Points in the Plane
by Eriol L. Lloyd 228

New NP-Hard and NP-Complete Polynomial and Integer Divisibility Problems
by David A. Plaisted 241

Lower Bounds for Natural Proof Systems (Preliminary Report)
by Dexter Kozen 254

On the Average Number of Registers Required for Evaluating Arithmetic Expressions
by P. Flajolet, J. C. Raoult, and J. Vuillemin

We write $M_n = B_n + \frac{1}{n} \left(\frac{2n}{n} \right) I_n + o(B_n)$

with $I_n = \int_0^{\infty} S_2((t/\sqrt{n})) e^{-t^2} \cdot H_3(t) dt$ (28)

Using Delange's formula (25),

$$U(x) = \int_0^x S_2((t/\sqrt{n})) dt = \frac{1}{2} x \log_2 x + x F(\log_2 x) + h(x),$$

we compute I_n by parts :

$$I_n = \frac{1}{\sqrt{n}} U(t/\sqrt{n}) e^{-t^2} H_3(t) \Big|_0^{\infty} - \frac{1}{\sqrt{n}} \int_0^{\infty} U(t/\sqrt{n}) e^{-t^2} H_4(t) dt.$$

The first term reduces to zero for at least two reasons, and :

$$I_n = \frac{1}{\sqrt{n}} \int_0^{\infty} \left[\frac{1}{2} (t/\sqrt{n}) \log_2(t/\sqrt{n}) + (t/\sqrt{n}) F(\log_2(t/\sqrt{n})) + h(t/\sqrt{n}) \right] e^{-t^2} H_4(t) dt.$$

Splitting in four :

$$\begin{aligned} I_{n_1} &= -\frac{1}{2} \int_0^{\infty} t \log_2 t H_4(t) e^{-t^2} dt - \frac{1}{4} \log_2 n \int_0^{\infty} t H_4(t) e^{-t^2} dt \\ &- \int_0^{\infty} t H_4(t) F(\log_2 n + \log_2 t) e^{-t^2} dt - \frac{1}{\sqrt{n}} \int_0^{\infty} h(t/\sqrt{n}) H_4(t) e^{-t^2} dt \\ &= -\frac{1}{2} I_1 - \frac{1}{4} \log_2 n I_2 - I_3 - \frac{1}{\sqrt{n}} I_4. \end{aligned}$$

Each integral is computed in turn.

$$\begin{aligned} I_1 &= \int_0^{\infty} t \log_2 t H_4(t) e^{-t^2} dt = -\frac{1}{\log 2} \int_0^{\infty} (\log t + 1) H_3(t) e^{-t^2} dt \\ &= -\frac{1}{\log 2} \int_0^{\infty} \log t (-8t^2 + 12t) e^{-t^2} dt - \frac{2}{\log 2} \\ &= -\frac{1}{\log 2} \int_0^{\infty} \log t^2 (-2t^2 + 3) e^{-t^2} dt - \frac{2}{\log 2} \\ &= -\frac{1}{\log 2} \int_0^{\infty} \log x (3-2x) e^{-x} dx - \frac{2}{\log 2} \\ &= -\frac{3}{\log 2} \int_0^{\infty} \log x e^{-x} dx + \frac{2}{\log 2} \int_0^{\infty} x \log x e^{-x} dx - \frac{2}{\log 2} \\ &= -\frac{1}{\log 2} \int_0^{\infty} \log x e^{-x} dx. \end{aligned}$$

This last integral is classical (see [Bu,33]) and its value is $-\gamma$, $\gamma \approx 0.577$ being Euler's constant.

Simple integrations by parts give $I_2 = -2$; as for I_4 , the function $h(t/\sqrt{n})$ ranges between 0 and 1, thus I_4 has some definite value and

$$\frac{1}{\sqrt{n}} I_4 = o(1).$$

There just remains $I_3 = K(\log_2 n)$, where

$$(29) \quad K(u) = \int_0^{\infty} t H_4(t) F(u + \log_2 t) e^{-t^2} dt$$

Since F is periodic with period 1 then so is K . The change of variable $x = u + \log_2 t$ shows that K is indefinitely differentiable.

We have thus proved :

Theorem : The average number $A_n = M_n/B_n$ satisfies

$$A_n = \log_4 n + D(\log_2 n) + o(1)$$

in which

$$D(u) = 1 - \frac{\gamma}{2 \log 2} + K(u),$$

$K(u)$ being the periodic function defined by (29).

Apart from the periodic term, one can, in principle, develop an asymptotic expansion as far as needed. H. Delange (private communication) has also computed the Fourier series of D :

$$D(t) = \sum_{k \in \mathbb{Z}} a_k e^{i2k\pi t}$$

and finds :

$$a_0 = \frac{1}{2} - \frac{\gamma + 2}{2 \log 2} + \log_2 \pi,$$

which is the mean value of $D(t)$, and for $k \neq 0$,

$$a_k = \frac{2k\pi i - \log 2}{\log 2} \Gamma\left(\frac{k\pi i}{\log 2}\right) \zeta\left(\frac{2k\pi i}{\log 2}\right)$$

where Γ and ζ are the classical gamma and Riemann's zeta functions.

The values of A_n have been machine computed for n ranging from 2 to 300 by means of formula (20). The results are plotted on figure 4, where the horizontal axis represents values of n in a logarithmic scale and the vertical axis represents values of $A_n - \log_4 n$.




Figure 4

The diagram shows an excellent agreement of A_n to $\log_4 n$ already for small values of n . It also shows the rapid convergence of $M_n - \log_4 n$ to the term $D(\log_2 n)$ centered around $a_0 = 0.292$, whose periodicity is apparent.

$$\frac{2k\pi i - \log 2}{\log 2} \Gamma\left(\frac{k\pi i}{\log 2}\right) \zeta\left(\frac{2k\pi i}{\log 2}\right)$$

Genesis of “Analytic Combinatorics”

Late 1970s / early 1980s: optimism and opportunity

Knuth volumes 1-3




Search for generality

Algorithms for the masses


Teaching and research in AofA



When will the book be available?



An alert: “best-seller **impatiemment attendu**”



From: Google Alerts <googlealerts-noreply@google.com>
Subject: Google Alert - sedgewick
Date: September 26, 2008 12:26:13 AM EDT
To: Bob Sedgewick

Google Blogs Alert for: **sedgewick**

[Best-seller **impatiemment attendu**](#)

By Michael Lugo(Michael Lugo)

I've seen citations to Flajolet and **Sedgewick's** book Analytic Combinatorics in things written as long ago as 1998 or so; Amazon.com says the book will be out on December 31, 2008, and the publisher, Cambridge University Press says ...

[God Plays Dice - http://godplaysdice.blogspot.com/](http://godplaysdice.blogspot.com/)

An alert: “best-seller **impatiemment attendu**”

God Plays Dice

A random walk through mathematics – mostly through the random part.

25 SEPTEMBER 2008

Best-seller **impatiemment attendu**

From “Quelques contributions au carrefour de la géométrie, de la combinatoire et des probabilités”, by [Nicolas Pouyanne](#):

“Dans leur best-seller **impatiemment attendu** [28], P. Flajolet et R. Sedgewick offrent un développement approfondi de la méthode symbolique en analyse combinatoire, auquel on pourra se référer.”

That is, “In their impatiently awaited best-seller [28], P. Flajolet and R. Sedgewick offer a detailed development of the symbolic method in combinatorial analysis, to which one will be able to refer.”

I don't laugh often while reading mathematics, but this made me laugh, because I am among those impatiently awaiting this book. I've seen citations to Flajolet and Sedgewick's book *Analytic Combinatorics* in things written as long ago as 1998 or so; [Amazon.com says the book is due December 31, 2008](#), and [the publisher, Cambridge University Press, says it will be published in December 2008](#). It can be downloaded from [Flajolet's web page](#).

Posted by Michael Lugo at 4:16 PM

Labels: [analytic combinatorics](#)

3 comments:

[Intrinsicallyknotted](#) said...

So does this count as “proof by impending publication”? Not “proof by vigorous handwaving”, but very amusing!

[September 25, 2008 7:10 PM](#)

[Michael Lugo](#) said...

Not really. I feel like “proof by impending publication” refers to things that aren't publicly available; Flajolet and Sedgewick's book is on the web. (I'm not sure if it will stay that way after publication. In any case, I'll buy a copy, because I spend enough time flipping back and forth between my PDF of it and other windows that having a paper copy would be more practical.)

[September 25, 2008 7:15 PM](#)


[David](#) said...

I downloaded it, but... woohoo, 800+ pages. I think the boss would be



Subscribe in a reader

1325 readers
BY FEEDBURNER



Blog Archive

▼ 2008 (343)

November (33)

... in their **impatiently awaited best-seller**, P. Flajolet and R. Sedgewick offer a detailed development of the symbolic method in combinatorial analysis, to which one will be able to refer.

Philadelphia, Pennsylvania,
United States

I'm a fourth-year PhD student in mathematics at the University of Pennsylvania. I study combinatorics and probability.

[View my complete profile](#)

“Impatiently awaited” on the web

Prince of Persia



plasma display




iPod




student visa



Berkeley Unix 4.4



Dan Brown novel



Heroes episode



“Impatiently awaited” in literature

*Prince Andrew listened to the account of the opening of the Council of State, which he had so **impatiently awaited** and to which he had attached such importance*

Leo Tolstoy, War and Peace

*Custer and his cavalry contingent **impatiently awaited** marching orders.*

Kingsley Bray, Crazy Horse: A Lakota Life

*This done, he **impatiently awaited** the return of his companions.*

Alexandre Dumas, The Count of Monte Cristo

*A handsome young fellow like you does not obtain long leaves of absence from his mistress, and we were **impatiently awaited** at Paris, were we not ?*

Alexandre Dumas, The Three Musketeers


*He **impatiently awaited** her husband's departure.*

Guy de Maupassant, Bel Ami

Digression: Separated at birth??



Impatiemment attendu



Last Sunday I sent the message that follows, then a reminder mid-week, but haven't received an ACK. Did you get it? Was it properly addressed?

Is there a pilot in the plane?

At full speed on the highway ... but in reverse gear?

Where are we????????

Did some e-mail message get lost? Please at least ACK having received the message below, first sent OCT 28, 2008.

Patience

Dear Phillippe,

Apologies for not acknowledging, but I have been working around the clock to get this book to press which it did this morning. All of your corrections have been incorporated and we are on schedule for a stock date of **28th November**.

Best Wishes,

[CUP staff member]



A calendar for November 2008. The days of the week are listed at the top: Sun, Mon, Tue, Wed, Thu, Fri, Sat. The dates are arranged in a grid. The date 28 is highlighted with a red circle.

November 2008						
Sun	Mon	Tue	Wed	Thu	Fri	Sat
26	27	28	29	30	31	1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29

From: david.tranah@gmail.com

Subject: Re: Is there a pilot in the plane?

Date: **November 28, 2008** 1:12:56 PM EST

To: Philippe.Flajolet@inria.fr

Cc: rs@CS.Princeton.EDU, dtranah@cambridge.org

Dear Philippe and Bob

Just a quick note to say that I am looking forward to meeting you both again next week at the 60th birthday colloquium, and that we are on schedule to have books on display for SODA in NYC in January and at the annual meeting of the American Mathematical Society in Washington, also in January.

best wishes

david

It's on the web. Why publish a book?

The New York Times

OP-ED CONTRIBUTOR

How to Publish Without Perishing

By JAMES GLEICK

Published: November 29, 2008

As a technology, the book is like a hammer. That is to say, it is perfect: a tool ideally suited to its task. Hammers can be tweaked and varied but will never go obsolete. Even when builders pound nails by the thousand with pneumatic nail guns, every household needs a hammer.

....Now even modest titles have been granted a gift of unlimited longevity.

What should an old-fashioned book publisher do with this gift? Forget about cost-cutting and the mass market. Don't aim for instant blockbuster successes. You won't win on quick distribution, and you won't win on price. Cyberspace has that covered.

Go back to an old-fashioned idea: that a book, printed in ink on durable paper, acid-free for longevity, is a thing of beauty. Make it as well as you can. People want to cherish it.



Analytic Combinatorics

Philippe Flajolet and
Robert Sedgewick

CAMBRIDGE